Introduction

Chairman Alexander, Ranking Member Murray and distinguished members of Committee. It is an honor to appear before you today to discuss the Department of Health and -19 pandemic. We are grateful for this opportunity to address how each of our agencies and offices are harnessing innovation to prevent, diagnose, and treat the novel coronavirus SARS-CoV-2.

COVID-

preparedness and response in the United States and abroad, and addressing infectious diseases like COVID-19 is central to our mission.

When there is an emerging pathogen like the SARS-CoV-2 virus, CDC expertise lies in our ability to study the new pathogen to understand how it is transmitted, and translate that knowledge into public health action. Since first learning of the cluster of cases in Wuhan, China, CDC has rapidly advanced the science around this new human pathogen, SARS-CoV-2. CDC has been both

protect Americans from infection. Currently, over 6,700 CDC employees have been engaged in

-19 response, and over 1,200 of these staff have been deployed to nearly 200 different locations in the United States and abroad. CDC staff have conducted rapid investigations of outbreaks that identified highest-risk priority populations and settings. Understanding specific population-level vulnerabilities and how infections spread in various types of settings has been instrumental in the development of guidance that will help keep the American people healthy and allow critical infrastructure services to be provided safely. For example, after data emerged that contrary to expectation, SARS-CoV-2 could be transmitted by people without symptoms, CDC recommended that people wear masks around others who do not

settings such as hair salons and childcare centers.

highest quality science has laid the foundation of what we know about COVID-19 and illuminated the way forward.

In addition to publishing our own scientific information on SARS-CoV-

to detect, respond, and prevent the spread of SARS-CoV-2 and to inform community mitigation strategies.

they need to address the COVID-19 pandemic. These jurisdictions are best positioned to understand the unique situation of each community, including the status of their public health infrastructure and workforce and its needs for enhancement. CDC is supporting STLT partners who are working to identify cases; conduct contact tracing; implement containment measures and mitigate spread of the virus in the community. CDC is working alongside these health departments to improve surveillance and reporting and enhance testing capacity. Together, STLT and CDC teams are responding to COVID-19 outbreaks in high-risk settings and implementing best practices to control the spread of the virus.

on managing disease outbreaks, CDC provides guidance and support for the development and implementation of effective containment and community mitigation strategies. The goal is for all jurisdictions to have robust public health systems, which include a contact tracing infrastructure that meets their unique needs. As of

increased monthly testing targets. These discussions and plans for action emphasize the need to serve disproportionately affected populations and include focused efforts for long-term care facilities, federally qualified health centers, American Indian/Alaska Native serving health facilities, and urban health facilities, among others.

CDC is working with STLT health departments to support forward-looking testing strategies that ensure that populations at higher risk, such as persons of color, have adequate access to testing. For example, CDC worked with the Health Resources and Services Administration (HRSA) and Federally Qualified Health Centers (FQHCs) to survey FQHCs and better understand the populations they are serving. Approximately 60 percent of responding

efforts when requesting an EUA. CDC took these steps to catalyze the development and validation of assays that can detect and differentiate SARS-CoV-2 from influenza by the commercial sector.

In March 2020, CDC and public health partners began seroprevalence surveys of community transmission of SARS-CoV-2. Seroprevalence surveys help identify infections that might be missed due to lack of symptoms or testing not being performed. Serology studies can also help determine risk factors associated with SARS-CoV-2 infection, including transmission in health care settings, and inform guidance and mitigation strategies. For example, CDC has published the results from one of the seroprevalence studies that used remnants of samples collected during routine clinical care. This was done in conjunction with two commercial companies and results suggested that greater than 10 times more SARS-CoV-2 infections occurred than the number of reported COVID-19 cases. Another study on healthcare personnel who routinely cared for COVID-19 patients found that 6 percent had evidence of previous SARS-CoV-2 infection. This study identified two factors potentially associated with SARS-CoV-2 infection among health care personnel: personal protective equipment (PPE) shortages and not wearing a mask while interacting with patients.

Data Collection, Analysis and Understanding of the Pandemic

CDC, in concert with HHS, continues to focus on data modernization efforts including expanding core data, informatics, and IT capacity; advancing interoperable systems and tools; strengthening and expanding collaboration with and support for partners and; coordinating data and IT investments and governance.

Accurate data are critical as we continue to assess the burden placed on the American healthcare system to inform reopening. CDC is leveraging all available surveillance systems, including influenza and viral respiratory disease systems, to monitor COVID-19 and protect disproportionately affected communities. These data collected by CDC help target critical COVID-19 interventions. In collaboration with STLT public health partners, CDC is committed to making data available to the public, while protecting individual privacy.

This crisis has highlighted the need to continue efforts to modernize the public health data systems that CDC and states rely on for accurate data. Public health data surveillance and analytical infrastructure modernization efforts started in FY 2020. Timely and accurate data are

essential as CDC and the nation work to understand the impact of COVID-19 on all Americans, particularly for populations at greater risk for severe illness, such as older Americans, those with chronic medical conditions, and some racial and ethnic minorities. Modernization efforts include support for the surveillance and data workforce, a key asset of the public health system. For example, CDC is working closely with our partners to help STLT health departments implement the Sara Alert system. Sara Alert is a standards-based, open source tool that automates the process of public health monitoring and reporting individuals exposed to, or infected with, COVID-19. To date, nine states and two territories, along with eight counties and one Tribal Council, have adopted the Sara Alert system. Almost 350,000 individuals have been monitored since April 8. During an average week, close to 80,000 individuals roll in and roll back off of monitoring through Sara Alert.

-based COVID-NET system monitors COVID-19 associated

identified as highest risk. For example, CDC is supporting local activities in Black and African American, Hispanic/Latino, American Indian and Alaska Native, and Asian American, Pacific Islander, and Native Hawaiian communities to deliver COVID-19 prevention messages and community mitigation strategies. CDC recently released a COVID-19 Health Equity Strategy (www.cdc.gov/coronavirus/2019-ncov/community/health-equity/cdc-strategy.html) that provides an evidence-based, comprehensive and coordinated framework for reducing COVID-19 disparities. The Strategy includes expanded plans for collecting and reporting timely, complete, representative, and relevant data on testing, incidence, vaccination, and severe outcomes among populations at highest risk. Additionally, CDC is working with existing program grantees, such as Racial and Ethnic Approaches to Community Health (REACH), and tribal grantees through a number of programs, to enhance outreach to populations at increased risk of complications from COVID-19. These broad-based community engagements and strategies are working with the aim of ensuring equitable access to testing, health care, and future COVID-19 vaccines.

American Indian and Alaska Native communities are some of the most affected by COVID-19. As of August 2020, CDC has provided $206.4 million to tribal nations, consortia, and organizations for responding to COVID-19 across tribal communities. This amount exceeds the minimum of $165 million directed by Congress through the Coronavirus Preparedness and Response Supplemental Appropriations Act and the CARES Act. CDC is using a multifaceted approach, guided by data, to allocate COVID-19 funding to tribal communities, enabling broad access to COVID-19 resources through a variety of direct and indirect supports.

Children

We are learning more about how COVID-19 impacts children every day. Although children are less likely than adults to develop severe illness when infected with SARS-CoV-2, household studies and outbreak investigations confirm that children can transmit the virus and often have the same or higher viral loads in their nasopharynx compared with adults. Though the mortality rate is low for children aged 18 years and younger, COVID-19 associated hospitalization rates increased among this age group during the summer. From March 1, 2020 to July 25, 2020, one in three hospitalized children was admitted to an intensive care unit.

CDC is committed to providing schools, teachers, staff, parents, and caregivers with information and guidance to help keep our children as safe and healthy as possible as schools

distribution and tracking. State and local health departments have conducted pandemic vaccination planning with immunization and preparedness funding from CDC for over a decade. Updating these vaccination response plans for implementation of COVID-19 vaccines will build readiness for timely administration when a vaccine becomes available. During August 2020, CDC completed in-person and virtual site visits to assess needs as vaccine planning intensifies. Lesson
jurisdictions to aid in the development of state-specific COVID-19 vaccination plans.

In addition, some state and local health departments utilized supplemental resources to build infrastructure that would address current COVID-19 response needs and incorporated planning for future phases. One example is in Chicago, where the health department has developed the Chi COVID Coach app to communicate directly with Chicago's residents who may be COVID-19 positive. The forward-thinking app, built by private sector companies, can be adapted throughout the course of the pandemic. It now allows users to register to receive a vaccine once they become available.

While it remains unclear how long the pandemic will last, COVID-19 activity will likely continue for some time. It is also unclear what impact the ongoing COVID-19 pandemic will have on health care and public health systems during the upcoming influenza season and beyond. Circulation of SARS-CoV-2 and influenza virus at the same time could place a tremendous burden on the health care system. Therefore, getting a seasonal flu vaccine is especially important. It is important that Americans have confidence in all vaccines. CDC will leverage its immunization program to help maintain high coverage in routine childhood immunizations, to

defeat COVID-19. As we work together to fight COVID-19 and end this pandemic, CDC is committed to its mission to protect all Americans from disease threats and to save lives, now and in the future.

National Institute of Allergy and Infectious Diseases

The National Institutes of Health (NIH) is the HHS agency leading the research response to COVID-19 and the novel coronavirus that causes the disease, SARS-CoV-2. Within NIH, NIAID is responsible for conducting and supporting research on emerging and re-emerging infectious diseases, including COVID-19.

NIAID responds rapidly to threats of emerging infectious diseases, by accelerating fundamental basic research efforts, engaging a domestic and international basic and clinical research infrastructure that can be quickly mobilized, and leveraging collaborative and highly productive partnerships with industry. NIAID also provides preclinical research resources to scientists in academia and private industry throughout the world to advance translational research on emerging and re-emerging infectious diseases. These research resources help bridge gaps in the product development pipeline, thereby lowering the scientific, technical, and financial risks incurred by product developers and incentivizing companies to partner with us in developing safe and effective countermeasures including vaccines, therapeutics, and diagnostics.

NIAID has a longstanding commitment to coronavirus research, including extensive efforts to combat two other serious diseases caused by coronaviruses: SARS and MERS. This research has enhanced our fundamental understanding of coronaviruses in general and provides a strong foundation for our accelerated efforts to address the specific challenge of COVID-19 by developing vaccines, therapeutics, and diagnostics.

Developing Vaccines and Monoclonal Antibodies to Prevent SARS-CoV-2 Infection and/or COVID-19

A safe and effective vaccine for SARS-CoV-2 will be essential to stopping the spread of infection, reducing rates of morbidity and mortality, and preventing future outbreaks.

NIAID recently established the COVID-19 Prevention Network (CoVPN) by leveraging four existing NIAID-

Consortium (IDCRC), and the AIDS Clinical Trials Group (ACTG), in partnership with the Department of Defense (DOD). The CoVPN aims to enroll thousands of volunteers in large-scale clinical trials testing a variety of investigational vaccines and monoclonal antibodies intended to protect people from SARS-CoV-2 infection and/or COVID-19. The CoVPN is a -private partnership led by HHS to invest in and coordinate the development, manufacture, and distribution of safe and effective COVID-19 vaccines, therapeutics, and diagnostics. The CoVPN is participating in harmonized protocols developed in collaboration with the Accelerating COVID-19 Therapeutic Interventions and Vaccines (ACTIV) public-private partnership, vaccine manufacturers, and Biomedical Advanced Research and Development Authority (BARDA). The network is participating in numerous trials at more than 100 clinical trial sites across the United States and internationally. The CoVPN has developed an extensive community engagement framework to reach out to the diverse communities most affected by COVID-19; to understand their interest in, and concerns about, research participation; and to partner with them to ensure their input is reflected in study

2020. The vaccine efficacy trial was the first to be initiated under OWS. The Coalition for Epidemic Preparedness Innovations (CEPI) funded the manufacture of the vaccine candidate for the Phase 1 trial, and BARDA is supporting advanced development of the candidate.

On August 31, 2020, NIAID announced the launch of a Phase 3 clinical trial with AstraZeneca of the AZD1222 COVID-19 vaccine candidate, which uses a chimpanzee adenovirus-vectored vaccine approach developed by researchers at the University of Oxford in collaboration with scien

On September 8, 2020, AstraZeneca paused enrollment in this and other studies of its candidate

settings, respectively. The ACTIV-2 and ACTIV-3 clinical trials utilize master protocols that allow for inclusion of additional investigational therapeutics as the trials continue. NIAID also is planning separate clinical trials to assess hyperimmune intravenous immunoglobulin for treatment of COVID-19 in both outpatients and hospitalized adults.

The National Heart, Lung, and Blood Institute (NHLBI) has established the Collaborating Network of Networks for Evaluating COVID-19 and Therapeutic Strategies (CONNECTS) to better understand the impact of COVID-19 on the heart, lungs, blood, and blood vessels, and to identify therapies that will slow or halt disease progression and speed recovery. CONNECTS will leverage existing NIH-funded clinical trial networks to conduct adaptive trials, in which

to identify potential SARS-CoV-2 therapeutics for further investigation. Other Institutes and Centers across NIH also are working concurrently with partners in academia and industry to pursue the development and testing of mAbs, antiviral, and anti-thrombotic drugs for potential treatment of COVID-19. NIAID, NHLBI, NCATS, the National Cancer Institute (NCI), the National Institute of Arthritis and Musculoskeletal and Skin Diseases, and the National Institute of Neurological Disorders and Stroke (NINDS) are all engaged in this critical effort.

NIH has established the COVID-19 Treatment Guidelines Panel, comprised of representatives of NIH and five other federal agencies along with representatives of nine professional organizations, academic experts, and treating physicians including providers from high COVID-19 incidence areas, and community representatives. On April 21, 2020, the panel issued the initial iteration of the COVID-19 treatment guidelines for clinicians. The guidelines provide recommendations regarding specific treatments and address considerations for special populations, including pregnant women and children. Based on a randomized controlled trial of the antiviral drug remdesivir compared to placebo, the Panel updated these treatment guidelines to recommend remdesivir for the treatment of COVID-19 in hospitalized patients who require supplemental oxygen with the caveat that due to insufficient data, the panel could not recommend for or against the use of remdesivir for those patients who require oxygen delivery through a high-flow device, noninvasive, ventilation, invasive mechanical ventilation, or ECMO. On June 25, 2020, based on a preliminary analysis of the data from the *Randomised Evaluation of COVID-19 Therapy* (RECOVERY) study sponsored by the University of Oxford, the treatment guidelines were updated again to recommend the glucocorticoid dexamethasone for the treatment of COVID-19 in hospitalized patients with severe disease requiring supplemental oxygen, including those on high flow oxygen or mechanical ventilation. Recently the treatment guidelines were updated to emphasize that potentially effective treatments for COVID-19 not be withheld from pregnant women. The guidelines are updated regularly as new evidence emerges.

In addition, the Pediatric Trials Network, funded by the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (NICHD), has incorporated testing of 12 drugs in its ongoing clinical trials that may prove helpful specifically for treating children with COVID-19 and/or multisystem inflammatory syndrome in children (MIS-C).

Enhancing Diagnosis and Understanding the Pathogenesis of COVID-19

NIH is supporting an HHS-wide effort to promote the development and commercialization of diagnostic tests to detect current SARS-CoV-2 infection. On April 29, 2020, NIH announced the Rapid Acceleration of Diagnostics (RADx) initiative, which is working to identify, support, and make innovative strategies for COVID-19 testing widely accessible, in collaboration with FDA, CDC, and BARDA. RADx is leveraging the Point-of-Care Technologies Research Network established by the National Institute of Biomedical Imaging and Bioengineering (NIBIB) to drive the development of new products by fall 2020. This $500 million initiative supports point-of-care and home-based diagnostic devices, as well as innovations that make current laboratory tests faster, more efficient, and more widely accessible. Innovators will be matched with technical, clinical, regulatory, business, and manufacturing experts to increase the odds of success. To date, NIH has awarded more than $370 million in Phase 2 contracts to 16 organizations for technology validation, clinical studies, scale-up, and manufacturing. These funds, combined with additional pending projects in the RADx pipeline, are projected to produce millions of new tests per day for a total of more than 6.5 million tests per day by the end of the year. In addition, NIAID is using CARES Act funds to support diverse SARS-CoV-2 diagnostic platforms including RT-PCR and enzyme-linked immunosorbent assays, and facilitating development of sensitive, specific, and rapid diagnostic tests by providing critical SARS-CoV-2 isolates and reagents to the developers of tests.

The RADx Underserved Populations (RADx-UP) initiative will augment the reach and power of technologies developed and enhanced through RADx by identifying and addressing implementation factors that present barriers to testing and follow-up in vulnerable populations. On June 12, 2020, NIH announced four new funding opportunities for community-engaged projects within RADx-UP. The goal of this is to better understand factors that have led to a disproportionate burden of the pandemic on underserved and vulnerable populations so that interventions can be implemented to decrease these disparities. Awards are expected to be made in late September or early October 2020.

NCI is coordinating with FDA and NIAID to assess the sensitivity and specificity of certain SARS-CoV-2 serological tests, which can detect antibodies indicative of a prior exposure to SARS-CoV-2. NCI and NIAID also are working to establish a collaborative national network to increase national capacity for high-
se nqBomaTl/F2 12 Tfseien BT/F2 12 TfseienBT/F2 12 TfseienBT/F2 12 TfseienBT 6729.8ud 6rOctober

In addition, they will conduct research to increase the understanding and application of those results and support related clinical efforts, including clinical trials of convalescent serum and the establishment of registries of tested subjects for seroprotection studies.

NIAID, NCI, and NHLBI, along with scientists from CDC, BARDA, FDA and DOD, convened the COVID-19 Serology Studies workshop to bring together over 300 scientists and clinicians from the federal government, industry, and academia on May 7, 2020. Participants discussed the role of serology testing in understanding and responding to the COVID-19 public health crisis and explored strategies to address key scientific opportunities and knowledge gaps in this emerging field. On July 14, 2020, a report of the conclusions and recommendations from the workshop was published in the journal *Immunity*. The group recommended that additional research is needed to determine whether, and to what extent, a positive antibody test means a person may be protected from reinfection with SARS-CoV-2. Additional research also is needed to determine the duration of protection. They also emphasized that serology tests should not be used as a stand-alone tool to make decisions about personal safety related to SARS-CoV-2 exposure until additional information about SARS-CoV-2 immunity is available.

NIAID, NCI, NCATS, and NIBIB also are partnering on a study to investigate whether adults in the United States without a confirmed history of infection with SARS-CoV-2 have antibodies to the virus, indicating prior infection. Public interest in participating was extremely robust and recruitment was significantly expanded by leveraging the NCATS Clinical and Translational Science Awards (CTSA) program. Enrollment is now complete. In addition, NIH is supporting COVID-19 natural history studies to understand the incidence of infection in specific populations, including children and their household contacts, and aspects of the clinical course of infection, including incidents of thrombosis, strokes, heart attacks, and other sequelae of infection. Some of these studies will examine the quality and durability of the immune response to SARS-CoV-2 and evaluate whether unique immune responses may be associated with clinical disease trajectories; this information may be leveraged to develop SARS-CoV-2 therapeutics or vaccines. Natural history studies also will inform our understanding of COVID-19 pathogenesis, including factors that may predict disease progression and help to identify individuals or groups at high risk.

In order to improve understanding of neurological consequences of SARS-CoV-2 and inform potential treatment strategies, NINDS is supporting a database that is collecting data on

disruption through targeted infection prevention and control measures. The indications for viral testing depend on the stage of the pandemic and the extent of community spread. In general, testing is indicated for diagnosis of those who are symptomatic or asymptomatic, tracing of those in contact with those who are infected, screening of specific employees (for example nursing home staff), and surveillance testing of those who are asymptomatic to achieve infection control and/or other public health objectives.

Repeated testing of a majority of the U.S. population is not feasible at this time, nor necessary to ensure a

Ensure sufficient supply of reagents to achieve testing goals in the context of point of care utilization and use of commercial referral labs;

Prioritize states with outbreaks or potential outbreaks, if needed, and;

Continue to expand the availability and use of point of care tests.

Support Public Health Isolation and Contact Tracing

A key function of testing is to support identification of SARS-CoV-2 of infected individuals, many of whom may be asymptomatic, in communities identified with outbreaks or emerging outbreaks. In respons

testing to increase baseline testing 2X-5X for short periods of time. Surge testing sites have been implemented in Miami, FL; Jacksonville, FL; Edinburg, TX; Yuma County, AZ; Pima County, AZ; Coconino, AZ; Phoenix, AZ; Atlanta, GA; Birmingham, AL; Cochise County, AZ; Mohave County, AZ; Yavapai County, AZ; Baton Rouge, LA; New Orleans, LA; Bakersfield, CA; Houston, TX; Harris County, TX; Clark County, NV; and Honolulu, HI, and these 19 sites have conducted approximately 290,000 tests. Surge testing is a supportive adjunctive activity; it cannot substitute for disciplined adherence to mitigation measures including masking, hygiene, social distancing, avoidance of indoor crowded areas and crowds, and protection of the vulnerable. These mitigation techniques, when combined with selective surge testing, have proven highly effective in reversing recent community outbreaks.

In order to support public health isolation and contact tracing, and to reduce turnaround time, the federal government has:

Provided massive surge testing to localities prioritized by the White House Coronavirus Taskforce, and agreed to by state and local officials;

Augmented testing, both baseline and surge, for FQHCs and retail sites;

Supported local testing efforts with surges of collection supplies and reagents;

Worked collaboratively to validate and promote EUAs for pooling across all laboratory platforms;

Worked collaboratively to validate and

Provided point-of-care testing to all nursing homes in America.

Diagnose COVID-19 Rapidly in Hospitalized Patients

Because there are now treatments authorized by the FDA for hospitalized patients with COVID-19, including remdesivir, convalescent plasma, and steroids, it is critical to diagnose patients as soon as possible. Currently, large commercial labs are prioritizing inpatient samples to ensure diagnosis within 24-36 hours. Our best information also suggests that the great majority of individual hospitals are able to meet this time frame for patients within their hospital systems.

Protect the Vulnerable

The elderly, particularly those in nursing homes, are much more likely to suffer serious consequences, including death, from COVID-19. In addition to the elderly, racial and ethnic minorities are also disproportionately affected.

To ensure that specimens are collected without overburdening the traditional health care system, and to ensure testing in the most vulnerable communiti(l)-.70 612 70.0047 8otionately affected.

To prevent further spread and deaths in nursing homes, CDC and the Centers for Medicare & Medicaid Services (CMS) recommended that nursing homes perform baseline testing of all residents and staff, followed by routine testing of residents and/or staff to reduce outbreaks, morbidity, and mortality, based on additional factors. CMS requires a regimen of staff and resident testing based on the degree of community spread.

To protect the vulnerable and to assist states in meeting these recommendations and requirements, on July 14, 2020, the Trump Administration announced that HHS would embark on a one-time procurement of rapid point-of-care testing instruments and tests to be distributed to nursing homes using the Defense Production Act. Through this aggressive action, nursing homes will be able to augment their current capacity for coronavirus testing, bolstering their response and helping to prevent the spread of SARS-CoV-2. This will facilitate baseline testing among nursing home residents and staff, and enable a pathway to conduct ongoing testing according to public health guidelines.

I am pleased to announce that all 13,850 initially eligible nursing homes have received one or more point-of-care (POC) instruments, and nearly 5 million tests. Following this initial distribution, we will facilitate nursing homes being able to reorder supplies via their normal commercial distribution channels. Additional billions of dollars in funding have been provided by HHS to support this effort.

Vulnerable populations in many underserved communities are suffering disproportionate health impacts resulting from COVID-19, including numbers of infections, hospitalizations, and deaths. As part of the HHS response to this crisis, on June 23, the HHS Office of Minority Health (OMH) announced the selection of the Morehouse School of Medicine as the awardee for a new $40 million initiative to fight COVID-19 in racial and ethnic minority, rural and socially vulnerable communities.

Morehouse School of Medicine has entered into a cooperative agreement with OMH to lead the initiative to coordinate a strategic network of national, state, territorial, tribal and local organizations to deliver COVID-19-related information to communities hardest hit by the pandemic. The three-year initiative will include the development and coordination of a strategic and structured network of national, state, territorial, and local public and community-based organizations that will help mitigate the impact of COVID-19 on racial and ethnic minorities as well as rural and socially vulnerable communities across the nation. The initiative also includes

a national multi-media outreach and education effort. One of the primary goals of these information dissemination efforts is to provide additional education and community-level information on resources to help fight the pandemic to those who need it most.

Support Safe Reopening of Schools and Businesses

While we must be prudent to protect those most vulnerable, we must also be mindful of the prolonged effects that school and business closures have on millions of children and parents. The efforts of the federal government to galvanize the testing infrastructure in the United States, and the efforts to reduce turnaround times, have provided communities with the resources they need to safely reopen schools and businesses.

Enable State Plans

To enable states to achieve the testing goals developed in coordination with the federal government, the federal government has worked with manufacturers to gain insight into diagnostic instrument install bases; procured and shipped collection supplies; and determined reagent inventory. The federal government then provided this information to states so they could better determine how to optimize their testing strategy. The federal government also purchased and allocated POC devices and over 2.3 million tests; developed, implemented, and facilitated community-based testing sites across the country; and provided significant guidance and technical assistance for state plans. The increase in the numbers of tests performed since early March is a direct reflection of these efforts.

to the public on August 10, and are available for viewing on the following website: https://www.hhs.gov/coronavirus/testing-plans/index.html

To ensure states meet their testing goals, the federal government procured FDA-authorized swabs and transport media, and is distributing these supplies to a single location in each state determined by the Governors offices. Starting in May and through September 11, the federal government has distributed over 95 million swabs and more than 77 million tubes of transport media.

Moving forward, jurisdictions should use the $10.25 billion provided to states, territories, and localities by the federal government to support the purchase of tests and related supplies, personnel for contact tracing, and reporting infrastructure, etc., for their jurisdictions, as needed to fulfill their approved testing plans.

Other Initiatives

In order to capture feedback and foster communication between federal officials and the private sector, HHS created the National Testing Implementation Forum (Forum). The Forum brings together representatives from key stakeholder groups to share information and provide input to federal leaders about SARS-CoV-2 testing. Members of the Forum provide their perspectives on how HHS can best identify and address end-to-end testing supply chain issues across commercial, public health, academic, and other sectors and define optimal testing in various settings (diagnostic, screening, surveillance, others). Members also provide input to improve technical assistance across the nation to prioritize testing among the vulnerable and underserved and create a sustainable diagnostics ecosystem that is sustainable and fully capable for future public health challenges. The first Forum meeting was held on July 30th and the principal topic of discussion was the testing supply chain. On August 13th the second meeting was held and surveillance and reopening strategies were discussed. The third forum, with the topic of engaging minority and underserved communities, was held on September 3rd.

On August 27th, the Administration announced that a $760 million contract was awarded to Abbott for the delivery of 150 million rapid BinaxNOW COVID-19 Ag Card point-of-care tests. This initiative will expand strategic testing in the United States. The Abbott BinaxNOW COVID-19 Ag Card, which recently received an EUA from the FDA, does not require

instrumentation and will deliver COVID-19 test results in 15 minutes or less. This test uses nasal swabs and can be easily deployed in many settings across the country.

United States Public Health Service Commissioned Corps

Since the early stages of the COVID-19 pandemic, the Corps has been an indispensable asset leveraged to address the public health needs of the nation in response to this crisis. The Corps is one of the eight uniformed services of the United States and the only uniformed service

industries we regulate with the tools and flexibility to do the same. Work has focused on facilitating the development and availability of medical countermeasures to diagnose, treat, and prevent COVID-19, surveilling the medical product and food supply chains for potential shortages or disruptions, and helping to mitigate such impacts, as necessary to protect the public health. This work is a key component of the efforts to address this pandemic and reopen the economy so Americans can get back to work and school.

Diagnostic Testing

This pandemic has created a demand for new tests that is unprecedented in both volume

for use in the U.S. provide sufficiently accurate and reliable results and helping to provide timely access to such tests.

Even prior to any diagnosed U.S. cases of COVID-19, FDA proactively reached out to developers to encourage and facilitate the development of tests and to offer assistance from the Agency. FDA has been proactive and supportive of test development by all interested parties to speed development and quickly authorize tests that the science supports. The Agency has worked

tests. However, flexibility never meant we would allow fraud. Unfortunately, FDA continues to see unscrupulous actors marketing fraudulent test kits and using the pandemic as an opportunity to take advantage of Americans. Some test developers have falsely claimed their tests are FDA-approved or authorized. Others have falsely claimed that their serology tests can diagnose COVID-19 or that they are authorized for at-home testing. FDA also became aware that a concerning number of commercial serology tests were performing poorly based on an independent evaluation by the NCI.

When we become aware of these issues, we have and will continue to take appropriate action against firms unlawfully marketing and selling their tests. FDA has and continues to issue Warning Letters and continues examining shipments of tests at ports of entry, the borders, and international mail centers, detaining and refusing fraudulent test kits.

To date, FDA has refused admission to more than 470 shipments of tests at the border, representing more than 460,000 tests overall, helping to prevent fraudulent tests from entering sellers of unapproved and unauthorized products removing false or misleading COVID-19 claims. FDA also has sent abuse complaints to online marketplaces and domain registrars for websites or listings that were offering to distribute fraudulent COVID-19 test kits for at-home testing. FDA has and will continue to take appropriate action against firms and individuals that place the public health at risk.

To further support efforts to ensure that patients and health care providers can depend on the results of COVID-19 tests, FDA has announced our participation in the COVID-19 Diagnostics Evidence Accelerator, a multi-stakeholder collaborative project to advance the development of diagnostics through the generation of real-world evidence. Organized by the Reagan-Udall Foundation for FDA in collaboration with Friends of Cancer Research, this initiative is designed to allow the community to analyze both diagnostic and clinical data in real time, which has the potential to contribute to the scientific evaluation of diagnostic tools and medical interventions for COVID-19.

Evidence generated by the Accelerator project is intended to be complementary to other studies that have been conducted or are underway as well as to provide actionable information about the prevalence of SARS-CoV-2 in specific populations and highlight individual risk factors for patients. This helps improve our understanding of the disease, allows us to tailor

public health interventions and strategies to mitigate risks for individuals and communities, and will help to stop the spread of SARS-CoV-2.

In addition, FDA also continues to work with NIH, the CDC, and BARDA regarding the

including antibody tests that are not the subject of an EUA or pre-EUA, as well as those that are under FDA review. Where appropriate, FDA is using NCI data to inform future decision making, such as whether to authorize the test, engage the test developer m 0 0 1 72.024 584.iduals be 0 1 o8Tf( )]TJETQ

time should be to support traditional FDA approval by conducting studies to directly evaluate the ability of the vaccine to protect humans from SARS-CoV-2 infection and/or disease.

In its interactions with vaccine developers, FDA provides sponsors with advice regarding the data needed to support the manufacturing, clinical development, and approval of vaccines, including such advice to those sponsors pursuing development of vaccines to prevent COVID-19. The size of clinical trials to evaluate the efficacy of COVID-19 vaccines will depend on a number of factors including the criteria for demonstrating safety, efficacy, and the incidence of COVID-19 in the population and areas where the trials are conducted. The guidance document conveys that FDA would expect that a COVID-19 vaccine would be at least 50 percent more effective than placebo in preventing COVID-19 or SARS-CoV-2 infection among the clinical trial participants. FDA anticipates that clinical trials to demonstrate vaccine efficacy would also be of sufficient size to provide an acceptable safety database. However, further pre-licensure safety evaluation may be needed if safety concerns arise during clinical development.

While FDA is committed to expediting this work, we will not cut corners in our decisions and are making clear through this guidance what data should be submitted to meet our regulatory standards. This is particularly important, as we know that some people are skeptical of efforts to develop a safe and effective COVID-19 vaccine. To help ensure an evaluation process that is as

uating the

safety and effectiveness of new vaccines, FDA will convene a meeting of our Vaccines and Related Biological Products Advisory Committee on October 22, 2020, to address the general development of COVID-19 vaccines. We stand ready to rapidly schedule additional meetings of this Committee after submission to discuss Biologic License Applications or request for an Emergency Use Authorization for COVID-19 vaccines.

It is clear that manufacturing and fill finish capacity will need to be scaled up on U.S. soil in order to have a safe and effective vaccine widely available in a timely manner. FDA is committed to working with sponsors by providing timely regulatory advice and technical assistance regarding manufacturing to help support such scale-up activities, including sponsors who may be proceeding at risk to scale-up manufacturing while clinical trials are being completed.

We have not lost sight of our responsibility to the American people to maintain our regulatory independence and ensure our decisions related to all medical products, including

COVID-19 vaccines, are based on science and the available data. This is a commitment that the American public can have confidence in and one that FDA will continue to uphold.

Therapeutic Development

Since the beginning of the COVID-19 pandemic, FDA has been working tirelessly to facilitate the development and availability of therapeutics for use by patients, physicians, and health systems as expeditiously and safely as possible. FDA announced on March 31, 2020, the creation of an emergency review and development program for possible therapies for COVID-

The Agency has supported the program by reassigning staff and working continuously to review requests from companies, scientists, and doctors who are working to develop therapies. Under CTAP, FDA is using every available authority and regulatory flexibility to facilitate the development of safe and effective products to treat patients with COVID-19.

Further, FDA is partnering with the NIH in its efforts to develop a national strategy for a coordinated research response to the pandemic. The ACTIV partnership developed a framework for prioritizing vaccine and drug candidates, streamlining related clinical trials, coordinating regulatory processes, and leveraging assets among all partners to rapidly respond to COVID-19 and future pandemics.

There are a variety of therapeutic products being evaluated, including antiviral drugs and immunotherapies that may be helpful in key areas, such as in reducing lung inflammation and improving lung function in COVID-19 patients. All this work is beginning to pay off. For example, we saw positive results of the National Institute of Allergy and Infectious Diseases (NIAID) trial of remdesivir in patients with severe COVID-19. On May 1, 2020, FDA issued an EUA for remdesivir for the treatment of suspected or laboratory-confirmed COVID-19 in adults and children hospitalized with severe disease. On August 28, 2020, FDA broadened the EUA for remdesivir to include all hospitalized patients for treatment of COVID-19, irrespective of their severity of disease.

Another potential approach for treatment is the use of antibody-rich products such as convalescent plasma and hyperimmune globulin. These investigational blood products are manufactured from plasma donated by people who have recovered from the SARS-CoV-2 virus, and such products are being studied to determine if they could shorten the length, or lessen the

treat hospitalized patients with COVID-19 when approved drugs are not available. The Agency has also published guidance to help applicants and manufacturers provide FDA with timely and informative notifications about changes in the production of certain drugs and biological products and urging the submission of these notifications, which may assist in our efforts to prevent or mitigate shortages of such products.

In addition, when we identify a shortage, we react swiftly to mitigate the impact to U.S. patients and health care professionals, and quickly share that information with the public. The Agency quickly identified the need for making hand sanitizers available as demand spiked, while also continuing our mission to ensuring these products remain safe for consumer use by removing adulterated products from the market. FDA has published and continues to update three guidance documents designed to help facilitate the production of alcohol-based hand sanitizer in non-traditional settings such as pharmacies or distilleries. The agency has initiated several enforcement initiatives and import alerts to stop adulterated and subpotent hand sanitizer

supply of medical devices to meet continuing unmet needs all over the country. FDA also provided instructions for importers to facilitate the import entry process for PPE and other devices, to help expand access to these products. To further support these efforts, FDA initiated biweekly virtual town hall meetings for those seeking and manufacturing respirators and other PPE to ask questions and discuss challenges they are facing.

We are also in close communication with our partners at U.S. Customs and Border Protection (CBP) to proactively identify and mitigate any potential backlogs of lawfully marketed medical products for COVID-19. FDA participates in HHS Supply Chain Advisory Group meetings, providing regulatory support and subject matter expertise to respond to questions concerning medical products identified by HHS, to facilitate the lawful entry and use of imported medical products coordinated through HHS, and to inform medical product supply chain discussions.

community have helped to accelerate patient access to critical devices. FDA appreciates Congress including provisions in the CARES Act for additional device shortages authority during or in advance of a declared public health emergency and looks forward to continuing to work with members of Congress to further expand these authorities, consistent with the FY 2021 Budget.

Inspections

Despite pausing on-site domestic and foreign routine surveillance inspections in March 2020 to safeguard the health and well-being of our staff, our investigators continued to conduct mission critical inspections both domestically and abroad and other activities to ensure FDA-regulated industries were meeting applicable FDA requirements. In July 2020, FDA resumed prioritized on-site domestic routine surveillance inspections. To arm our investigators with the most reliable and accurate information, the FDA developed a rating system to assist us in determining when and where it is safest to conduct prioritized domestic inspections. The COVID-19 Advisory Rating system (COVID-19 Advisory Level) uses real-time data to qualitatively assess the number of COVID-19 cases in a local area based on state and national data. We have also made the Advisory Level data available to our state partners who carry out inspections of FDA- We will continue

and criminal enforcement actions, where appropriate. In March 2020, FDA launched Operation